

# THE COMBINATORICS OF MOTZKIN POLYOMINOES

JEAN-LUC BARIL, SERGEY KIRGIZOV, JOSÉ L. RAMÍREZ, AND DIEGO VILLAMIZAR

**ABSTRACT.** A word  $w = w_1 \cdots w_n$  over the set of positive integers is a Motzkin word whenever  $w_1 = 1$ ,  $1 \leq w_k \leq w_{k-1} + 1$ , and  $w_{k-1} \neq w_k$  for  $k = 2, \dots, n$ . It can be associated to a  $n$ -column Motzkin polyomino whose  $i$ -th column contains  $w_i$  cells, and all columns are bottom-justified. We reveal bijective connections between Motzkin paths, restricted Catalan words, primitive Lukasiewicz paths, and Motzkin polyominoes. Using the aforementioned bijections together with classical one-to-one correspondence with Dyck paths avoiding  $UDUs$ , we provide generating functions with respect to the length, area, semiperimeter, value of the last symbol, and number of interior points of Motzkin polyominoes. We give asymptotics and closed-form expressions for the total area, total semiperimeter, sum of the last symbol values, and total number of interior points over all Motzkin polyominoes of a given length. We also present and prove an engaging trinomial relation concerning the number of cells lying at different levels and first terms of the expanded  $(1 + x + x^2)^n$ .

## 1. INTRODUCTION

In the literature, Dyck and Motzkin paths play a crucial role in many problems from combinatorial theory (see [19] and the references therein). Remember that a *Dyck path* of semilength  $n$  is a lattice path of  $\mathbb{Z} \times \mathbb{Z}$  running from  $(0, 0)$  to  $(2n, 0)$  that never passes below the  $x$ -axis and whose permitted steps are  $U = (1, 1)$  and  $D = (1, -1)$ . A *Motzkin path* of length  $n$  is a lattice path in the first quadrant made of  $n$  steps  $U$ ,  $D$ , and  $F = (1, 0)$ , from the origin to the point  $(n, 0)$ . These paths are enumerated by the well-known Catalan and Motzkin numbers (respectively, [A000108](#) and [A001006](#) in the On-line Encyclopedia of Integer Sequences [29]).

These sequences have a similar behavior since, almost always, they appear together, and a large number of similar types of combinatorial classes are enumerated by these numbers. For Catalan numbers, we refer to the catalog of Catalan combinatorial classes compiled by Stanley [30]. Similarly, there are several combinatorial classes where the Motzkin family arises. For example, the rooted plane trees are a classical Catalan family, but if we add loops, yields a Motzkin family [14]. Additional examples of Motzkin families can be found in [4, 8] and the references therein.

In this paper, we focus on the class of Motzkin words introduced by Mansour and Ramírez in [25]. Specifically, a word  $w = w_1 w_2 \cdots w_n$  of length  $n$  over the set of positive integers is called a *Motzkin word* whenever  $w_1 = 1$ ,  $1 \leq w_k \leq w_{k-1} + 1$ , and  $w_{k-1} \neq w_k$  for  $k = 2, \dots, n$ . The empty word  $\epsilon$  is the only one word of length 0. For  $n \geq 0$ , let  $\mathcal{M}_n$  denote the set of Motzkin words of length  $n$ . For example,

$$\mathcal{M}_5 = \{12121, 12123, 12312, 12321, 12323, 12341, 12342, 12343, 12345\}.$$

---

*Date:* June 23, 2024.

*2010 Mathematics Subject Classification.* 05A15, 05A19.

*Key words and phrases.* Motzkin word, generating function, bijections.

In [25], the authors prove that the cardinality of the set  $\mathcal{M}_n$  is given by the Motzkin number  $m_{n-1}$ , where  $m_n$  is defined by the combinatorial sum (see [8, 14])

$$m_n = \frac{1}{n+1} \sum_{i \geq 0} \binom{n+1}{i} \binom{n+1-i}{i+1}, \quad n \geq 0.$$

The number  $m_n$  corresponds to the  $n$ -th term of the generating function

$$M(x) := \sum_{n \geq 0} m_n x^n = \frac{1 - x - \sqrt{1 - 2x - 3x^2}}{2x^2},$$

and the first few values of the Motzkin numbers are

$$1, \quad 1, \quad 2, \quad 4, \quad 9, \quad 21, \quad 51, \quad 127, \quad 323, \quad 835, \quad 2188, \dots$$

For all  $1 \leq k \leq n$ , we denote by  $\mathcal{M}_{n,k}$  the set of the Motzkin words of length  $n$  whose last symbol is  $k$ . Let  $m(n, k)$  denote the number of Motzkin words in  $\mathcal{M}_{n,k}$ . It is clear that  $m_n = \sum_{k \geq 1} m(n, k)$ . Moreover, in [25] the authors proved that

$$(1) \quad m(n, k) = \frac{k}{n} \sum_{j=k}^n (-1)^{n-j} \binom{n}{j} \binom{2j-k-1}{j-1}.$$

A Motzkin word  $w = w_1 \cdots w_n$  can also be viewed as a polyomino (also called bargraph) whose  $i$ -th column contains  $w_i$  cells for  $1 \leq i \leq n$ , and where all columns are bottom-justified (i.e. the bottom-most cells of all of its columns are in the same row). The polyomino associated to a Motzkin word of length  $n$  is called a *Motzkin polyomino* of length  $n$ . In Figure 1, we show all Motzkin polyominoes of length 5.




FIGURE 1. Motzkin words of length 5 and their associated Motzkin polyominoes.

In this work, we focus on the class of Motzkin polyominoes, and we investigate enumerative problems about several statistics on these objects.

Let  $w$  be a Motzkin word and  $P(w)$  its associated polyomino. We denote by  $\mathbf{area}(w)$  the number of cells (or the *area*) of  $P(w)$ , which also is the sum of all  $w_i$  for  $1 \leq i \leq n$ . The *semiperimeter* of  $P(w)$ , denoted  $\mathbf{sper}(w)$ , is half of the *perimeter* of  $P(w)$ , while the perimeter of  $P(w)$  is the number of cell borders that do not touch another cell of  $P(w)$ . An *interior-vertex* of  $P(w)$  is a point that belongs to exactly four cells of  $P(w)$ . We denote by  $\mathbf{inter}(w)$  the number of interior points of  $P(w)$ . For instance, if  $w = 12341$ , then  $\mathbf{area}(w) = 11$ ,  $\mathbf{sper}(w) = 9$ , and  $\mathbf{inter}(w) = 3$ . We refer to [18] for a historical review on polyominoes, and to [23] for the definitions of many statistical and enumerative methods over polyominoes.

Additionally, analogous results are known for bargraphs [12, 9], compositions [10], set partitions [21, 22], Catalan words [13, 26, 24], inversion sequences [1], and words [20, 11].

**Motivation and outline of the paper.** The motivation of this work is to provide enumerative results on Motzkin words according to several parameters (area, semiperimeter, and number of interior points) defined on the associated polyominoes. Except for the first section where we exhibit bijections between Motzkin words and other classical combinatorial classes counted by the Motzkin numbers, all other results are obtained algebraically by considering functional equations for multivariate generating functions. In Section 2 we give three bijections between Motzkin words and three known combinatorial classes counted by the Motzkin numbers. In Section 3, we focus on the two statistics of the area and the semiperimeter. We provide the trivariate generating function, where the coefficient of  $x^n p^k q^\ell$  is the number of Motzkin polyominoes  $P(w)$  of length  $n$ , and satisfying  $\text{sper}(w) = k$  and  $\text{area}(w) = \ell$ . From this, we deduce the generating function for the total sum of the last symbol over all Motzkin words of length  $n$ , and we give asymptotic approximation for the coefficient of  $x^n$ . In Section 4, we focus on the semiperimeter statistic. We give the generating function for nonempty Motzkin polyominoes with respect to the length and the semiperimeter. We exhibit a bijection between Motzkin polyominoes of length  $n$  with semiperimeter  $2n - k$ , and Motzkin paths of length  $n - 1$  with exactly  $k$  up steps. We deduce a closed-form expression for the total semiperimeter over all Motzkin polyominoes of length  $n$ . In Section 5, we make a similar study as in Section 4 by considering the area instead of the semiperimeter. Finally, Section 6 is dedicated to the statistic of the number of interior points of the Motzkin polyominoes, and we exhibit a closed-form expression for the total number of interior points over all Motzkin polyominoes of length  $n$ .

## 2. LINKS BETWEEN MOTZKIN WORDS AND OTHER MOTZKIN CLASSES

Below, we present three bijections between Motzkin words (or equivalently Motzkin polyominoes) and three known combinatorial classes counted by the Motzkin numbers.

First, let us recursively define the bijection  $\psi$  between  $\mathcal{M}_n$  and the set of Motzkin paths of length  $n - 1$ , i.e. lattice paths in the first quadrant, starting at the origin, ending on the  $x$ -axis, and made of  $(n - 1)$  steps  $F = (1, 0)$ ,  $U = (1, 1)$ , and  $D = (1, -1)$ :

$$\psi(w) = \begin{cases} \epsilon & \text{if } w = 1, \\ F\psi(u) & \text{if } w = 1(1 + u), \text{ where } u \neq \epsilon, \\ U\psi(u)D\psi(v) & \text{if } w = 1(1 + u)v, \text{ where } u, v \neq \epsilon, \end{cases}$$

where  $u$  and  $v$  are Motzkin words and  $(1 + u)$  corresponds to the word obtained from  $u$  by increasing by one all letters of  $u$ . For instance, we have

$$\begin{aligned} \psi(12123453412) &= U\psi(1)D\psi(123453412) = UDU\psi(123423)D\psi(12) \\ &= UDUF\psi(12312)DF\psi(1) = UDUFU\psi(12)D\psi(12)DF = UDUFUFD\psi(12)DF. \end{aligned}$$

In Figure 2 we show the corresponding Motzkin path.

$$\psi(12123453412) = UDUFUFD FDF$$




FIGURE 2. Bijection between Motzkin paths and Motzkin words.

On the other hand, in a previous paper [6], the authors study the avoidance of patterns in Catalan words of length  $n$ , i.e. words  $w = w_1w_2 \cdots w_n$  satisfying  $w_1 = 0$  and  $0 \leq w_i \leq w_{i-1} + 1$ . They focus on Catalan words of length  $n$  avoiding the pattern  $(\geq, \geq)$ , that is those that do not contain occurrences  $w_i \geq w_{i+1} \geq w_{i+2}$ , or equivalently those avoiding the consecutive patterns 100, 000, 110, and 210. They prove algebraically that these words are counted by the Motzkin number  $m_n$ . Below, we give a bijection  $\phi$  establishing a link between this class of Catalan words and  $\mathcal{M}_n$ .

Let  $w$  be a Catalan word avoiding  $(\geq, \geq)$ , then  $\phi$  is recursively defined as follows.

$$\phi(w) = \begin{cases} 1 & \text{if } w = \epsilon, \\ 1(1 + \phi(u)) & \text{if } w = 1(1 + u), \\ 1(1 + \phi(\bar{u}))\phi(v) & \text{if } w = 1(1 + u)v, \quad u, v \neq \epsilon, \\ 1(1 + \phi(u))1 & \text{if } w = 11(1 + u), \\ 1(1 + \phi(\bar{u}))\phi(v)1 & \text{if } w = 11(1 + u)v, \quad u, v \neq \epsilon, \end{cases}$$

where  $u$  and  $v$  are some Catalan words, and  $\bar{u}$  is obtained from  $u$  by deleting the last symbol, and where  $(1 + u)$  corresponds to the word obtained from  $u$  by increasing by one all letters of  $u$ . For instance, we have

$$\phi(1123231231) = 1\phi(121)\phi(1231)1 = 12323123121.$$

We leave it as an exercise to check that  $\phi$  is a bijection.

**Link with primitive Łukasiewicz paths.** A *Łukasiewicz path* of length  $n \geq 0$  is a lattice path in the first quadrant of the  $xy$ -plane starting at the origin  $(0, 0)$ , ending on the  $x$ -axis, and consisting of  $n$  steps lying in the set  $\{(1, k) : k \geq -1\}$ . It is well-known [5, 16, 30] that these paths are enumerated by the Catalan numbers  $c_n = \frac{1}{n+1} \binom{2n}{n}$ . A *primitive* Łukasiewicz path is one with exactly one return on the  $x$ -axis (necessarily at the end). The empty path is not primitive. Let  $\mathcal{L}_n$  be the family of primitive Łukasiewicz paths of length  $n$  without horizontal steps.

There is a bijection between the sets  $\mathcal{M}_n$  and  $\mathcal{L}_{n+1}$ . Indeed, for  $1 \leq i \leq n$  the  $i$ -th entry of the Motzkin word is exactly the  $y$ -coordinate of the final point of the  $i$ -th step of the corresponding Łukasiewicz path (see Figure 3), and the avoidance of horizontal steps becomes from the avoidance of two consecutive identical entries in the Motzkin word.




FIGURE 3. Bijection between Motzkin words and primitive Łukasiewicz paths.

### 3. AREA AND SEMIPERIMETER STATISTICS

In this section, we study the area and semiperimeter statistics on Motzkin words (or equivalently Motzkin polyominoes). Recall that, for all  $1 \leq i \leq n$ ,  $\mathcal{M}_{n,i}$  is the set of Motzkin words of length  $n$  whose last symbol is  $i$ . We define the generating functions

$$A_i(x; p, q) := \sum_{n \geq 1} x^n \sum_{w \in \mathcal{M}_{n,i}} p^{\text{sper}(w)} q^{\text{area}(w)}.$$

and

$$A(x; p, q; v) := \sum_{i \geq 1} A_i(x; p, q) v^{i-1}.$$

**Theorem 3.1.** *The generating function  $A(x; p, q; v)$  satisfies the functional equation*

$$(2) \quad A(x; p, q; v) = p^2 q x + \frac{p q x}{1 - q v} A(x; p, q; 1) + \left( p^2 q^2 x v - \frac{p q x}{1 - q v} \right) A(x; p, q; q v).$$

*Proof.* Let  $w$  be a Motzkin word in  $\mathcal{M}_n$ . We distinguish two cases: (i)  $w \in \mathcal{M}_{n,1}$  and (ii)  $w \in \mathcal{M}_{n,i}$  for  $i \geq 2$ .

Case (i). The last symbol of  $w$  is 1. Hence, we have either  $w = 1$  or  $w = w'1$  where  $w' \in \mathcal{M}_{n-1,j}$ , with  $n \geq 2$  and  $j \geq 2$ . See Figure 4 for a graphical representation of these two cases. The generating function for this case is given by

$$A_1(x; p, q) = p^2 q x + p q x \sum_{j \geq 2} A_j(x; p, q).$$

From this relation it is clear that

$$(3) \quad A_1(x; p, q) = p^2 q x + p q x (A(x; p, q; 1) - A_1(x; p, q)).$$





FIGURE 4. Motzkin polyominoes whose last column is of height 1.

Case (ii). The last symbol of  $w$  is at least 2. From the definition of Motzkin word we have the decomposition  $w = w'i$ , with  $w' \in \mathcal{M}_{n-1,j}$ , and with either  $j = i - 1 \geq 1$  or  $j > i \geq 2$ . See Figure 5 for a graphical representation of these two cases.

FIGURE 5. Motzkin polyominoes whose last column is of height  $\geq 2$ .

In terms of generating functions, we deduce the functional equations:

$$(4) \quad A_i(x; p, q) = p^2 q^i x A_{i-1}(x; p, q) + pq^i x \sum_{j \geq i+1} A_j(x; p, q), \quad \text{for } i \geq 2.$$

By multiplying equation (4) by  $v^{i-1}$  and summing over  $i \geq 2$ , we obtain

$$\begin{aligned} A(x; p, q; v) - A_1(x; p, q) &= \sum_{i \geq 2} p^2 q^i x A_{i-1}(x; p, q) v^{i-1} + \sum_{i \geq 2} pq^i x \sum_{j \geq i+1} A_j(x; p, q) v^{i-1} \\ &= p^2 q^2 x v A(x; p, q; qv) + px \sum_{j \geq 3} A_j(x; p, q) \sum_{i=2}^{j-1} q^i v^{i-1} \\ &= p^2 q^2 x v A(x; p, q; qv) + px \sum_{j \geq 3} A_j(x; p, q) \frac{q^2 v - q^j v^{j-1}}{1 - qv} \\ &= p^2 q^2 x v A(x; p, q; qv) + \frac{pq^2 x v}{1 - qv} (A(x; p, q; 1) - A_1(x; p, q) - A_2(x; p, q)) \\ &\quad - \frac{pqx}{1 - qv} (A(x; p, q; qv) - A_1(x; p, q) - qv A_2(x; p, q)) \\ &= p^2 q^2 x v A(x; p, q; qv) + \frac{pq^2 x v}{1 - qv} (A(x; p, q; 1) - A_1(x; p, q)) \\ &\quad - \frac{pqx}{1 - qv} (A(x; p, q; qv) - A_1(x; p, q)). \end{aligned}$$

Simplifying the last expression using (3), we obtain the desired result.  $\square$

Note that when  $p = q = 1$ , we have

$$\left(1 - xv + \frac{x}{1-v}\right) A(x; 1, 1; v) = x + \frac{x}{1-v} A(x; 1, 1; 1).$$

The generating function  $A(x; 1, 1; 1)$  corresponds to the generating function of the nonempty Motzkin words, so  $A(x; 1, 1; 1) = xM(x)$ . Thus, we deduce the following.

**Corollary 3.2.** *The generating function for the number of nonempty Motzkin words with respect to the length and the value of the last symbol is*

$$(5) \quad A(x; 1, 1; v) \cdot v = \frac{xv(1-v) + x^2vM(x)}{1-v+x-vx+v^2x}.$$

The first terms of the series expansion are

$$xv + v^2x^2 + (v^3 + v)x^3 + (v^4 + 2v^2 + v)x^4 + (v^5 + 3v^3 + 2v^2 + 3v)x^5 + \\ (v^6 + 4v^4 + 3v^3 + 7v^2 + 6v)x^6 + (v^7 + 5v^5 + 4v^4 + 12v^3 + 14v^2 + 15v)x^7 + O(x^8).$$

There are three polyominoes of length 5 ending with a column of height three (see Figure 1). By differentiating  $A(x; 1, 1; v) \cdot v$  at  $v = 1$ , we deduce:

**Corollary 3.3.** *The generating function for the total sum  $g_n$  of the last symbol in all Motzkin words of length  $n$  is*

$$\frac{1 - x - 2x^2 - \sqrt{1 - 2x - 3x^2}}{2x^2}.$$

An asymptotic approximation for the coefficient  $g_n$  of  $x^n$  is given by

$$\frac{3\sqrt{3} \left(\frac{1}{n}\right)^{\frac{3}{2}} 3^n}{2\sqrt{\pi}}.$$

The asymptotic approximation is easily obtained with a singularity analysis (see [15, 27]).

The first terms of  $g_n$ ,  $1 \leq n \leq 10$ , are

$$1, \quad 2, \quad 4, \quad 9, \quad 21, \quad 51, \quad 127, \quad 323, \quad 835, \quad 2188, \dots$$

This sequence corresponds to the Motzkin numbers [A001006](#) in the OEIS [29]. Finally, the expected value of the last symbol is  $m_n/m_{n-1}$ , and an asymptotic is 3.

Notice that we can retrieve Corollary 3.3 as follows. From a Motzkin word  $w \in \mathcal{M}_{n-1,k}$ , we can construct  $k$  Motzkin words  $w' = wx \in \mathcal{M}_{n,x}$  by adding a letter  $x \in [1, k-1] \setminus \{k\}$ . Conversely, any Motzkin word of length  $n$  can be uniquely constructed with this process. This means that  $\sum_{k=1}^{n-1} km(n-1, k) = m_n$  which gives a simple combinatorial interpretation of Corollary 3.3.

#### 4. THE SEMIPERIMETER STATISTIC

In this section we study the semiperimeter statistic on Motzkin polyominoes. By (2) with  $q = 1$ , we obtain

$$(6) \quad \left(1 - p^2xv + \frac{px}{1-v}\right) A(x; p, 1; v) = p^2x + \frac{px}{1-v} A(x; p, 1; 1).$$

In order to compute  $S(x, p) := A(x; p, 1; 1)$ , we use the kernel method (see [2, 28]). The method consists in cancelling the coefficient of  $A(x; p, 1; v)$  by taking the small root  $v_0$  (the ones going to 0 for  $x \sim 0$ ) of  $1 - p^2 x v_0 + \frac{px}{1-v_0}$ , namely

$$v_0 = \frac{1 + p^2 x - \sqrt{1 - 2p^2 x - 4p^3 x^2 + p^4 x^2}}{2p^2 x}.$$

When the factor on the left-hand side equals zero (and thus satisfies the equation), the right-hand side also equals zero and so the solution  $v_0$  can be used on the right-hand side to obtain  $A(x; p, 1, 1)$  (see Theorem 4.1).

**Theorem 4.1.** *The generating function for the number of nonempty Motzkin polyominoes according to the length and the semiperimeter is given by*

$$S(x, p) = \frac{1 - p^2 x - \sqrt{1 - 2p^2 x - 4p^3 x^2 + p^4 x^2}}{2px}.$$

The first terms of the series expansion of  $S(x, p)$  are

$$p^2 x + p^4 x^2 + (p^5 + p^6) x^3 + (3p^7 + p^8) x^4 + (2p^8 + 6p^9 + p^{10}) x^5 + O(x^6).$$

Figure 6 yields the 6 Motzkin polyominoes of length 5 and semiperimeter 9.




FIGURE 6. Motzkin polyominoes of length 5 and semiperimeter 9.

**Theorem 4.2.** *The map  $\psi$  (defined in Section 2) induces a bijection between the Motzkin polyominoes (or equivalently words) in  $\mathcal{M}_n$  of semiperimeter  $2n - k$  for  $k = 1, 2, \dots, \lfloor \frac{n-1}{2} \rfloor$ , and the Motzkin paths of length  $n - 1$  with exactly  $k$  up steps.*

*Proof.* Let us prove the statement by induction on  $n$ . The case  $n = 0$  is trivial since  $\psi(1) = \epsilon$ . Let us assume the statement is true for  $m \leq n$ , and let us prove it for  $n + 1$ . Let  $w \in \mathcal{M}_{n+1}$  with semiperimeter  $2(n + 1) - k$ . We denote by  $\text{up}(M)$  the number of up-steps of the Motzkin path  $M$ .

If  $w = 1(1 + u)$ ,  $u \neq \epsilon$ , then we have  $\text{sper}(u) = 2n - k$ ,  $\psi(w) = F\psi(u)$ , and  $\text{up}(\psi(w)) = \text{up}(\psi(u))$ . Using the induction hypothesis on  $u$ , we have  $\text{up}(\psi(u)) = k$ , and then  $\text{up}(\psi(w)) = k$  as expected.

If  $w = 1(1 + u)v$ ,  $u, v \neq \epsilon$ , then we have  $\text{sper}(u) = s_1$  and  $\text{sper}(v) = s_2$  with  $s_1 + s_2 + 1 = 2(n + 1) - k$ ,  $\psi(w) = U\psi(u)D\psi(v)$ , and  $\text{up}(\psi(w)) = 1 + \text{up}(\psi(u)) + \text{up}(\psi(v))$ . Using the induction hypothesis on  $u$  and  $v$ , we have  $\text{up}(\psi(u)) = 2|u| - s_1$ , and  $\text{up}(\psi(v)) = 2|v| - s_2$ , where  $|u|$  is the length of  $u$ . Hence, this implies that  $\text{up}(\psi(w)) = 1 + 2|u| - s_1 + 2|v| - s_2 = 2 + 2|u| + 2|v| - 2(n + 1) + k = k$ , as expected.

Considering these two cases, the induction is completed.  $\square$



Using relation (6), we deduce the following.

**Theorem 4.3.** *The generating function for the number of nonempty Motzkin polyominoes according to the length, the semiperimeter, and the value of the last letter is given by*

$$A(x; p, 1; v) = \frac{1 + p^2(x - 2vx) - \sqrt{1 - 2p^2x - 4p^3x^2 + p^4x^2}}{2(1 + px + p^2v^2x - v(1 + p^2x))}.$$

The first terms of the series expansion of  $A(x; p, 1; v)$  are

$$p^2x + p^4vx^2 + (p^5 + p^6v^2)x^3 + (p^7 + 2p^7v + p^8v^3)x^4 + (2p^8 + p^9 + 2p^9v + 3p^9v^2 + p^{10}v^4)x^5 + O(x^6).$$

From Figure 6 we can verify that there are 1, 2, and 3 Motzkin polyominoes of length 5 and semiperimeter 9, whose value of the last symbol (height of last column) is 1, 2, and 3, respectively.

Let  $s(n, i)$  denote the sum of the semiperimeters of all Motzkin polyominoes of length  $n$  whose last column has height  $i$ . The first few values are

$$[s(n, i)]_{n, i \geq 1} = \begin{pmatrix} 2 & 0 & 0 & 0 & 0 & 0 & \cdots \\ 0 & 4 & 0 & 0 & 0 & 0 & \cdots \\ 5 & 0 & 6 & 0 & 0 & 0 & \cdots \\ 7 & 14 & 0 & 8 & 0 & 0 & \cdots \\ 25 & 18 & 27 & 0 & 10 & 0 & \cdots \\ 61 & 72 & 33 & 44 & 0 & 12 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

Notice that from the decomposition given in Figure 5, we have for  $n \geq 2$  and  $2 \leq i \leq n$ ,

$$s(n, i) = s(n-1, i-1) + 2m(n-1, i-1) + \sum_{j=i+1}^{n-1} (s(n-1, j) + m(n-1, j)),$$

where  $m(n, i)$  is given in (1). If we consider the difference  $s(n, i) - s(n, i-1)$ , then for  $n \geq 2$  and  $3 \leq i \leq n$ , we obtain the recurrence relation

$$s(n, i) = s(n, i-1) + s(n-1, i-1) - s(n-1, i-2) - s(n-1, i) + 2(m(n-1, i-1) - m(n-1, i-2)) - m(n-1, i).$$

Let  $s(n)$  be the total semiperimeter over all Motzkin polyominoes of length  $n$ . By Corollary 4.3 at  $v = 1$ , we deduce:

**Corollary 4.4.** *The generating function of the sequence  $s(n)$  is given by*

$$\frac{1 + x^2 - (1+x)\sqrt{1-2x-3x^2}}{2x\sqrt{1-2x-3x^2}}.$$

An asymptotic approximation of  $s(n)$  is given by

$$\frac{5\sqrt{3}\sqrt{\frac{1}{n}}3^n}{6\sqrt{\pi}}.$$

The first few values for  $1 \leq n \leq 10$  are

$$2, \quad 4, \quad 11, \quad 29, \quad 80, \quad 222, \quad 624, \quad 1766, \quad 5030, \quad 14396, \dots$$

This sequence does not appear in the OEIS. The expected value of the semiperimeter is  $5n/3$ . In Corollary 4.5 we give an explicit relation for the sequence  $s(n)$  using the *central trinomial coefficient*  $T_n$ , that is the coefficient of  $x^n$  in the expansion  $(x^2 + x + 1)^n$ . From the multinomial theorem we see that

$$T_n = \sum_{k=0}^n \binom{n}{k} \binom{n-k}{k}.$$

Moreover, the generating function of the central trinomial coefficients is given by

$$(7) \quad T(x) := \sum_{n \geq 0} T_n x^n = \frac{1}{\sqrt{1 - 2x - 3x^2}}.$$

**Corollary 4.5.** *The total semiperimeter over all Motzkin polyominoes of length  $n$  is given by*

$$s(n) = T_n + 2T_{n-1} - m_{n-1},$$

where  $m_n$  is the  $n$ -th Motzkin number.

*Proof.* By differentiating  $S(x, p)$  at  $p = 1$ , Theorem 4.1 gives

$$\begin{aligned} \frac{\partial}{\partial p} S(x, p)|_{p=1} &= \frac{1 + x^2 - (1 + x)\sqrt{1 - 2x - 3x^2}}{2x\sqrt{1 - 2x - 3x^2}} \\ &= -1 + T(x) + 2xT(x) - xM(x). \end{aligned}$$

Comparing the  $n$ -th coefficient we obtain the desired result.  $\square$

We leave as an open question to find a combinatorial interpretation of the equation obtained in Corollary 4.5.

## 5. THE AREA STATISTIC

The goal of this section is to analyze the area statistic on Motzkin polyominoes. By (2) with  $p = 1$  we obtain

$$(8) \quad A(x; 1, q; v) = qx + \frac{qx}{1 - qv} A(x; 1, q; 1) - \frac{qx(1 - qv + q^2v^2)}{1 - qv} A(x; 1, q; qv).$$

By iterating this equation an infinite number of times (here we assume  $|x| < 1$  or  $|q| < 1$ ), we obtain

$$A(x; 1, q; v) = \sum_{j \geq 1} (-1)^{j-1} q^j x^j \left( 1 + \frac{A(x; 1, q; 1)}{1 - q^j v} \right) \prod_{i=1}^{j-1} \frac{1 - q^i v + q^{2i} v^2}{1 - q^i v}.$$

By setting  $v = 1$ , and solving for  $A(x; 1, q; 1)$ , we can state the following result.

**Theorem 5.1.** *The generating function  $U(x, q) := A(x; 1, q; 1)$  for the number of nonempty Motzkin polyominoes according to the length and the area is given by*

$$U(x, q) = \frac{\sum_{j \geq 1} (-1)^{j-1} q^j x^j \prod_{i=1}^{j-1} \frac{1 - q^i + q^{2i}}{1 - q^i}}{1 - \sum_{j \geq 1} (-1)^{j-1} \frac{q^j x^j}{1 - q^j} \prod_{i=1}^{j-1} \frac{1 - q^i + q^{2i}}{1 - q^i}}.$$

The first terms of the series expansion of  $U(x, q)$  are

$$qx + q^3x^2 + (q^4 + q^6)x^3 + (q^6 + q^7 + q^8 + q^{10})x^4 + \\ + (q^7 + 3q^9 + 2q^{11} + q^{12} + q^{13} + q^{15})x^5 + O(x^6).$$

We refer to Figure 7 for an illustration of the polyominoes of length 5.




FIGURE 7. Motzkin polyominoes of length 5 and their weighted area.

**Theorem 5.2.** *The generating function for the number of Motzkin polyominoes, including the empty word, according to the length and area is given by the infinite continued fraction*

$$1 + U(x, q) = \frac{1}{1 - \frac{qx}{1 + qx - \frac{(1 + qx)q^2x}{1 + q^2x - \frac{(1 + q^2x)q^3x}{\ddots}}}}.$$

*Proof.* First note that the Motzkin words of length  $n$  are in bijection with Dyck paths of length  $2n$  avoiding the consecutive subword  $UDU$ . Indeed, given a Dyck path avoiding  $UDU$ , we associate a Motzkin word formed by the  $y$ -coordinate of each final point of the up steps. For example, in Figure 8 we show the Dyck path associated to the Motzkin word  $123212343 \in \mathcal{M}_9$ . We denote by  $\mathcal{D}$  the set of Dyck paths avoiding  $UDU$ .




FIGURE 8. Dyck path associated to the Motzkin word 123212343.

From this bijection, it is clear that the area of a Motzkin polyomino is equivalent to count the area of a Dyck path avoiding the subword  $UDU$  (the area of a Dyck path is the area of the region bounded by the path and the  $x$ -axis). Given a Dyck path  $P$  in  $\mathcal{D}$  from the first return decomposition it can be one of the following options:  $\epsilon, UD, UP'DP''$ , where  $P', P'' \in \mathcal{D}$  and

$P' \neq \epsilon$ . Let  $J(x, q)$  be the bivariate generating function for the Dyck paths in  $\mathcal{D}$  according to the length and the area. From the decomposition we obtain the functional equation

$$J(x, q) = 1 + xq + xq(J(qx, q) - 1)J(x, q).$$

Therefore, we deduce

$$J(x, q) = \frac{1 + xq}{1 - xq(J(qx, q) - 1)}.$$

Iterating this expression we obtain the desired result.  $\square$

Let  $u(n, i)$  denote the total area of the Motzkin polyominoes of length  $n$  that end with a column of height  $i$ . The first few values are

$$[u(n, i)]_{n, i \geq 1} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \cdots \\ 0 & 3 & 0 & 0 & 0 & 0 & \cdots \\ 4 & 0 & 6 & 0 & 0 & 0 & \cdots \\ 7 & 14 & 0 & 10 & 0 & 0 & \cdots \\ 27 & 21 & 33 & 0 & 15 & 0 & \cdots \\ 75 & 89 & 45 & 64 & 0 & 21 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

From the decomposition given in Figure 5, we have for  $n \geq 2$  and  $2 \leq i \leq n$ ,

$$u(n, i) = u(n-1, i-1) + i \cdot m(n-1, i-1) + \sum_{j=i+1}^{n-1} (u(n-1, j) + i \cdot m(n-1, j)),$$

where  $m(n, i)$  is given in (1). If we consider the difference  $u(n, i) - u(n, i-1)$ , then for  $n \geq 2$  and  $3 \leq i \leq n$ , we obtain the recurrence relation

$$\begin{aligned} u(n, i) &= u(n, i-1) + u(n-1, i-1) - u(n-1, i) - u(n-1, i-2) \\ &\quad + i \cdot m(n-1, i-1) - (i-1)m(n-1, i-2) - i \cdot m(n-1, i) + \sum_{j=i}^{n-1} m(n-1, j). \end{aligned}$$

Let  $u(n)$  be the total area of over all Motzkin polyominoes of length  $n$ . The first few values are

$$1, \quad 3, \quad 10, \quad 31, \quad 96, \quad 294, \quad 897, \quad 2727, \quad 8272, \quad 25048, \dots$$

This sequence corresponds to the sequence [A055217](#) in the OEIS [29]. Recently, Goy and Shattuck [17] gave a combinatorial interpretation for this sequence by using marked Motzkin paths. Our interpretation is different and probably new. In the following theorem we give an interesting combinatorial formula to calculate the sequence  $u(n)$ .

**Theorem 5.3.** *The total area over all Motzkin polyominoes of length  $n$  is given by*

$$u(n) = \frac{1}{2} (3^n - T_n) = \frac{1}{2} \left( 3^n - \sum_{k=0}^n \binom{n}{k} \binom{n-k}{k} \right),$$

and an asymptotic is  $3^n/2$ .

*Proof.* Let  $B(x; v) = \frac{\partial}{\partial q} A(x; 1, q; v) |_{q=1}$ . Then by differentiating (8) with respect to  $q$ , we obtain

$$B(x; v) = x + \frac{x^2 M(x)}{(1-v)^2} + \frac{x}{1-v} B(x; 1) - \frac{x(1-2v+4v^2-2v^3)}{(1-v)^2} A(x; 1, 1; v) - \frac{x(1-v+v^2)}{1-v} (B(x; v) + v \frac{\partial}{\partial v} A(x; 1, 1; v)).$$

By using (5), we obtain

$$(9) \quad \frac{\partial}{\partial v} A(x; 1, 1; v) = \frac{\partial}{\partial v} \left( \frac{x(1-v) + x^2 M(x)}{1-v+x-vx+v^2x} \right)$$

$$(10) \quad = \frac{1-2vx-x^2-2vx^2+2v^2x^2-(1+x-2vx)\sqrt{1-2x-3x^2}}{2(1-v+x-vx+v^2x)^2}.$$

Therefore

$$\begin{aligned} \frac{(1-v+x-vx+v^2x)^3}{1-v} B(x; v) &= x(1-v+x-vx+v^2x)^2 + \frac{x^2(1-v+x-vx+v^2x)^2 M(x)}{(1-v)^2} \\ &\quad + \frac{x(1-v+x-vx+v^2x)^2}{1-v} B(x; 1) \\ &\quad - \frac{x(1-2v+4v^2-2v^3)(1-v+x-vx+v^2x)}{(1-v)^2} (x(1-v) + x^2 M(x)) \\ &\quad - \frac{xv(1-v+v^2)}{2(1-v)} \left( 1-2vx-x^2-2vx^2+2v^2x^2-(1+x-2vx)\sqrt{1-2x-3x^2} \right). \end{aligned}$$

By twice differentiating this equation with respect to  $v$  and taking  $v = xM(x) + 1$ , we obtain that

$$B(x, 1) = \frac{1+x-\sqrt{1-2x-3x^2}}{2-4x-6x^2} = \frac{1}{2} \left( \frac{1}{1-3x} - \frac{1}{\sqrt{1-2x-3x^2}} \right).$$

Comparing the  $n$ -th coefficient we obtain the desired result.  $\square$

**5.1. Link with trinomial coefficients.** The sequence  $u(n)$  ([A055217](#)) corresponds to the sum of the first  $n$  coefficients of  $(1+x+x^2)^n$ . Enigmatically, the monomials from the first part of the expansion of  $(1+x+x^2)^n$  can be literally written on the cells of all Motzkin polyominoes of size  $n$  in a simple one-to-one manner. Term  $x^k$  goes onto a cell of height  $n-k$ , see Figure 9. This subsection is devoted to explain this fact, which is easy to state, but not easy to prove. Theorem 5.3 follows from the results of this section if we replace  $x$  by 1, consider the symmetry of coefficients of  $(1+x+x^2)^n$ , and subtract the central trinomial coefficient.

Let  $T(n, i)$  be the  $i$ -th coefficient in the expansion of  $(1+x+x^2)^n$ . It is not difficult to prove that

$$(11) \quad T(n, i) = T(n-1, i) + T(n-1, i-1) + T(n-1, i-2), \quad 0 \leq i \leq n-2,$$

and  $T(n, n-1) = T_{n-1} + T(n-1, n-2) + T(n-1, n-3)$ , where  $T_n$  is the central trinomial coefficient, that is  $T_n = T(n, n)$ . We also have  $T(n, n-1) = n|\mathcal{M}_n| = nm_{n-1}$ , see [A005717](#)




FIGURE 9. Correspondence between Motzkin polyominoes of size  $n$  and first  $n$  terms of the expansion of  $(1 + x + x^2)^n$ .

in the OEIS for more information. If  $i > n - 1$  or  $i < 0$  we define  $T(n, i) = 0$ , moreover  $T(1, 1) = 1$ . The first few values of this array are

$$[T(n, i)]_{n \geq 1, i \geq 0} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 1 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 1 & 3 & 6 & 0 & 0 & 0 & 0 & 0 & \dots \\ 1 & 4 & 10 & 16 & 0 & 0 & 0 & 0 & \dots \\ 1 & 5 & 15 & 30 & 45 & 0 & 0 & 0 & \dots \\ 1 & 6 & 21 & 50 & 90 & 126 & 0 & 0 & \dots \\ 1 & 7 & 28 & 77 & 161 & 266 & 357 & 0 & \dots \\ 1 & 8 & 36 & 112 & 266 & 504 & 784 & 1016 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

Let  $h(n, i)$  be the total number of cells of height  $i$  in  $\mathcal{M}_n$ . From Figure 9, we have  $h(5, 1) = 45, h(5, 2) = 30, h(5, 3) = 15, h(5, 4) = 5, h(5, 5) = 1$ . Notice that these numbers are related to the first 5 coefficients of the expansion of

$$(1 + x + x^2)^5 = \mathbf{1} + \mathbf{5x} + \mathbf{15x^2} + \mathbf{30x^3} + \mathbf{45x^4} + 51x^5 + 45x^6 + 30x^7 + 15x^8 + 5x^9 + x^{10}.$$

Let  $w$  be a Motzkin word. We denote by  $\mathbf{h}_i(w)$  the number of cells of height  $i$  in the Motzkin polyomino associated with  $w$ . We introduce the following generating functions

$$H_i(x, q) := 1 + \sum_{w \in \mathcal{M}} x^{|w|} q^{\mathbf{h}_i(w)}$$

and

$$B_i(x) := \left. \frac{\partial H_i(x, q)}{\partial q} \right|_{q=1}.$$

From the definition it is clear that  $[x^n]B_i(x) = h(n, i)$ .

**Theorem 5.4.** *For  $i \geq 2$ , we have*

$$H_i(x, q) = \frac{1 + x}{1 - (H_{i-1}(x, q) - x)},$$

and  $H_1(x, q) = 1 + qxM(xq)$ , where  $M(x)$  is the generating function of the Motzkin numbers.

*Proof.* Motzkin words of length  $n$  are in bijection with Dyck paths of length  $2n$  avoiding the subword  $UDU$  (the  $i$ -th entry of the word  $w$  corresponds to the  $y$ -coordinate of the endpoint of the  $i$ -th up-step of its corresponding Dyck path). From this bijection it is clear that

$$H_1(x, q) = 1 + xq + xq(H_1(x, q) - 1)H_1(x, q).$$

Solving this equation, we obtain that  $H_1(x, q) = 1 + qxM(xq)$ . Analogously, from the decomposition given in the proof of Theorem 5.2 we obtain the functional equation

$$H_i(x, q) = 1 + x + x(H_{i-1}(x, q) - 1)H_i(x, q), \quad i \geq 2. \quad \square$$

For example, for  $i = 3$  we have the expression

$$\begin{aligned} H_3(x, q) &= \frac{2 - x - q(2 + x^2) + x\sqrt{(1 + qx)(1 - 3qx)}}{2(1 - q - x + q(1 - x)x)} \\ &= 1 + x + x^2 + (1 + q)x^3 + (1 + 2q + q^2)x^4 + (1 + 3q + 3q^2 + 2q^3)x^5 + O(x^6). \end{aligned}$$

Figure 10 yields the Motzkin polyominoes of length 5 and the cells of height 3.




FIGURE 10. Motzkin polyominoes of length 5 and the cells of height 3.

**Proposition 5.5.** *For  $i \geq 1$ , we have*

$$B_i(x) = \frac{x^i M^i(x)}{1 - x - 2x^2 M(x)},$$

where  $M(x)$  is the generating function of the Motzkin numbers.

*Proof.* We proceed by induction on  $i$ . The identity clearly holds for  $i = 1$ ; see Theorem 5.4. Now suppose that the result is true for  $i - 1$ . We prove it for  $i$ :

$$\begin{aligned} B_i(x) &= \left. \frac{\partial H_i(x, q)}{\partial q} \right|_{q=1} = \left. \frac{\partial}{\partial q} \left( \frac{1+x}{1 - (H_{i-1}(x, q) - x)} \right) \right|_{q=1} = \left. \frac{\partial}{\partial q} \left( \frac{1}{1 - \frac{x}{1-x} H_{i-1}(x, q)} \right) \right|_{q=1} \\ &= \frac{\frac{x}{1+x} \frac{x^{i-1} M^{i-1}(x)}{1-x-2x^2 M(x)}}{\left( 1 - \frac{x}{1+x} (1+xM(x)) \right)^2} = \frac{(1+x)x^i M^{i-1}(x)}{(1-x^2 M(x))^2} = \frac{x^i M^i(x)}{1-x-2x^2 M(x)}. \end{aligned}$$

□

A *Motzkin walk* of length  $n$  is a Motzkin path prefix without the condition that never passes below the  $x$ -axis (see for instance [3]). A Motzkin walk ending on  $x$ -axis is also known as *Grand Motzkin path*. Let  $g(n, i)$  be the number of Motzkin walks of length  $n$  ending at height  $i$ . It is clear that  $g(n, i) = g(n-1, i-1) + g(n-1, i) + g(n-1, i+1)$ . We define the generating function

$$G_i(x) = \sum_{n \geq 0} g(n, i) x^n.$$

**Theorem 5.6.** *For all  $i \geq 0$ ,  $G_i(x) = B_i(x)$ . Moreover, For  $n > 0, i > 1$ , we have*

$$h(n, i) = h(n-1, i-1) + h(n-1, i) + h(n-1, i+1),$$

$$h(n, 1) = n \cdot [x^{n-1}]M(x).$$

*Proof.* Let  $P$  be a Motzkin walk ending at height  $i$ . It can be decomposed as  $GUM_1UM_2U \cdots UM_i$ , where  $G$  is a Motzkin walk ending at height zero and  $M_j$  is a Motzkin path for all  $1 \leq j \leq i$ . Therefore, we have the generating function  $G_i(x) = x^i G_0(x) M^i(x)$ , where  $M(x)$  is the generating function of the Motzkin numbers. Notice that  $G_0(x) = 1 + xG_0(x) + 2x^2 M(x)G_0(x)$ , because a nonempty Motzkin walk ending at height zero is either  $FG'_0$ , or  $UMDG'_0$ , or  $DMUG'_0$  where  $G'_0$  is a Motzkin walk ending at height zero,  $M$  is a Motzkin path (ending on the  $x$ -axis) and  $\bar{M}$  is a Motzkin walk obtained from a Motzkin path (ending on the  $x$ -axis) after exchanging  $U$  and  $D$ . So, we have  $G_0(x) = 1/(1-x-2x^2 M(x))$ . Therefore,

$$G_i(x) = \frac{x^i M^i(x)}{1-x-2x^2 M(x)} = B_i(x).$$

Now, it is clear that the sequences  $h(n, i)$  and  $g(n, i)$  are the same, therefore  $h(n, i)$  satisfies the desired recurrence relation. □

From Theorem 5.6 and (11), it is possible to verify that the sequence  $T(n, i)$  and  $h(n, n-i)$  satisfy the same recurrence relation with the same initial values, therefore they are the same.

**Theorem 5.7.** *The number of cells of height  $n-i$  ( $0 \leq i < n$ ) in all Motzkin polyominoes of length  $n$  is given by the trinomial coefficient  $T(n, i)$ .*

Now we can give an alternative proof of the combinatorial identity given in the Theorem 5.3.

*Second proof of Theorem 5.3.* Notice that  $\sum_{i=0}^{n-1} T(n, i) = u(n)$ . But the trinomial coefficients are palindromic, that is  $T(n, i) = T(n, 2n-i)$ , then

$$\sum_{i=0}^{2n} T(n, i) = 2u(n) - T(n, n) = 2u(n) - T_n.$$



Notice that we have to subtract the central trinomial coefficient. But it is clear that this sum is  $3^n$ , therefore  $3^n = 2u(n) - T_n$ .

We end this section by asking the following (open) questions: can we obtain an interpretation moving down a level to binomial coefficients? Can we obtain a generalization of this result by using tetranomial coefficients (i.e. coefficients of the polynomials  $(1 + x + x^2 + x^3)^n$ ), and for which polyomino classes?

## 6. THE INTERIOR POINTS STATISTIC

In this section, we study the statistic of the number of interior points on Motzkin polyominoes. As in the previous section, for all  $1 \leq i \leq n$ ,  $\mathcal{M}_{n,i}$  is the set of the Motzkin words of length  $n$  whose last symbol is  $i$ , and we define the generating functions

$$A_i(x; q) := \sum_{n \geq 1} x^n \sum_{w \in \mathcal{M}_{n,i}} q^{\text{inter}(w)}.$$

and

$$A(x; q; v) := \sum_{i \geq 1} A_i(x; q) v^{i-1}.$$

**Theorem 6.1.** *The generating function  $A(x; q; v)$  is given by*

$$A(x; q; v) = \sum_{j \geq 1} x^j \left( 1 + A(x; q; 1) \frac{1}{1 - q^j v} \right) \prod_{i=1}^{j-1} \left( xq^{i-1}v - \frac{x}{1 - q^i v} \right).$$

*Proof.* According to the decomposition given in Figures 4 and 5, we obtain the relations

$$\begin{aligned} A_1(x; q) &= x + x \sum_{j \geq 2} A_j(x; q), \\ A_i(x; q) &= xq^{i-2}A_{i-1}(x; q) + xq^{i-1} \sum_{j \geq i+1} A_j(x; q). \end{aligned}$$

By multiplying the last equation by  $v^{i-1}$  and summing over  $i \geq 2$ , we obtain the functional equation

$$\begin{aligned} A(x; q; v) - A_1(x; q) &= \frac{xqv}{1 - qv} A(x; q; 1) + \left( xv - \frac{x}{1 - qv} \right) A(x; q; qv) \\ &\quad - \left( \frac{xqv}{1 - qv} - \frac{x}{1 - qv} \right) A_1(x; q). \end{aligned}$$

Simplifying this expression we obtain the equation

$$(12) \quad A(x; q; v) = x + \frac{x}{1 - qv} A(x; q; 1) + \left( xv - \frac{x}{1 - qv} \right) A(x; q; qv).$$

By iterating the last equation an infinite number of times (here we assume  $|x| < 1$  or  $|q| < 1$ ), we obtain the desired result.  $\square$

By setting  $v = 1$  in Theorem 6.1, and solving for  $A(x; q; 1)$  we can state the following result.

**Corollary 6.2.** *The generating function  $H(x, q) := A(x; q; 1)$  for the number of nonempty Motzkin polyominoes according to the length and the number of interior points is given by*

$$H(x, q) = \frac{\sum_{j \geq 1} x^j \prod_{i=1}^{j-1} \left( q^{i-1} - \frac{1}{1-q^i} \right)}{1 - \sum_{j \geq 1} \frac{x^j}{1-q^j} \prod_{i=1}^{j-1} \left( q^{i-1} - \frac{1}{1-q^i} \right)}.$$

The first terms of the series expansion of  $H(x, q)$  are

$$H(x, q) = x + x^2 + (1 + q)x^3 + (1 + q + q^2 + q^3)x^4 + (1 + 2q + q^2 + 2q^3 + q^4 + q^5 + q^6)x^5 + O(x^6).$$

We refer to Figure 11 for an illustration of the polyominoes of length 5.




FIGURE 11. Motzkin polyominoes of length 5 and their weighted interior points.

**Corollary 6.3.** *The generating function of the number of interior points over all Motzkin polyominoes of length  $n$  is*

$$\frac{2 - 3x - 5x^2 - (2 - x - 2x^2)\sqrt{1 - 2x - 3x^2}}{2x(1 + x)(1 - 3x)},$$

and an asymptotic for the  $n$ -th coefficient of the series expansion is  $3^n/2$ . The expected value of the number of interior points is  $\sqrt{\frac{\pi}{3}}n^{3/2}$ .

*Proof.* Let  $B(x; v) = \frac{\partial}{\partial q} A(x; q; v) |_{q=1}$ . Then by differentiating (12) with respect to  $q$ , we obtain

$$\begin{aligned} B(x; v) &= \frac{vx^2 M(x)}{(1-v)^2} + \frac{x}{1-v} B(x; 1) \\ &\quad - \frac{xv}{(1-v)^2} A(x; 1, v) + \left( xv - \frac{x}{1-v} \right) (B(x; v) + v \frac{\partial}{\partial v} A(x; 1; v)). \end{aligned}$$

From (9), we know that

$$\frac{\partial}{\partial v} A(x; 1; v) = \frac{1 - 2vx - x^2 - 2vx^2 + 2v^2x^2 - (1 + x - 2vx)\sqrt{1 - 2x - 3x^2}}{2(1 - v + x - vx + v^2x)^2}.$$

Therefore

$$\begin{aligned} \frac{(1-v+x-vx+v^2x)^3}{1-v} B(x;v) &= \frac{x(1-v+x-vx+v^2x)^2}{1-v} B(x;1) \\ &\quad + \frac{v}{2} (1-x-vx-3x^2+2vx^2-v^2x^2-x^3+3vx^3-3v^2x^3+2v^3x^3 \\ &\quad - (1-xv-x^2+vx^2-v^2x^2)\sqrt{1-2x-3x^2}). \end{aligned}$$

By twice differentiating this equation with respect to  $v$  and taking  $v = xM(x) + 1$ , we obtain that

$$B(x,1) = \frac{2-3x-5x^2-(2-x-2x^2)\sqrt{1-2x-3x^2}}{2x(1+x)(1-3x)}.$$

Comparing the  $n$ -th coefficient we obtain the desired result.  $\square$

As an application of Pick's theorem (cf. [7, pp. 40]), one may establish a relation between the area, semiperimeter, and number of interior points of a Motzkin polyomino. Pick's theorem says that the area of a simple polygon with integer vertex coordinates is equal to  $I + B/2 - 1$ , where  $I$  is the number of interior points and  $B$  is the number of points lying on the boundary. Let  $\text{int}(n)$  be the sum of the interior points over all Motzkin polyominoes of length  $n$ .

**Corollary 6.4.** *The number of interior points over all Motzkin polyominoes of length  $n$  is*

$$\frac{1}{2} (3^n - 3T_n) - 2T_{n-1} + 2m_{n-1}.$$

*Proof.* From Pick's theorem we can obtain the relation

$$u(n) = \text{int}(n) + s(n) - m_{n-1}.$$

Now the identity follows from Corollary 4.5 and Theorem 5.3.  $\square$

The first few values of the sequence  $\text{int}(n)$  for  $n \geq 3$  are

$$1, \quad 6, \quad 25, \quad 93, \quad 324, \quad 1088, \quad 3565, \quad 11487, \dots$$

This sequence does not appear in the OEIS.

**Acknowledgments.** The authors are grateful to the referees for the detailed comments and corrections that helped us improve the paper. José L. Ramírez was partially supported by Universidad Nacional de Colombia. The third author would like to thank the Laboratoire d'Informatique de Bourgogne for the warm hospitality during his visit at Université de Bourgogne where part of this work was done. This work was supported in part by ANR-22-CE48-0002 funded by l'Agence Nationale de la Recherche and by the project ANER ARTICO funded by Bourgogne-Franche-Comté region (France).

## REFERENCES

- [1] M. Archibald, A. Blecher, and A. Knopfmacher. Parameters in inversion sequences, *Math. Slovaca* **73** (3) (2023), 551–564.
- [2] C. Banderier, M. Bousquet-Mélou, A. Denise, P. Flajolet, D. Gardy, and D. Gouyou-Beauchamps. Generating functions for generating trees, *Discrete Math.* **246** (2002), 29–55.
- [3] C. Banderier and P. Flajolet. Basic analytic combinatorics of directed lattice paths, *Theoret. Comput. Sci.* **281** (2002), 37–80.
- [4] J.-L. Baril and J.-M. Pallo. Motzkin subposets and Motzkin geodesics in Tamari lattices, *Inform. Process. Lett.* **114** (2014), 31–37.

- [5] J.-L. Baril and H. Prodinger. Enumeration of partial Łukasiewicz paths, *Enumer. Combin. Appl.* **3**(1) (2023), S2R2.
- [6] J.-L. Baril and J. L. Ramírez. Descent distribution on Catalan words avoiding ordered pairs of relations, *Adv. in Applied Math.* **149** (2023), 102551.
- [7] M. Beck and S. Robins. *Computing the Continuous Discretely: Integer Point Enumeration in Polyhedra*. Undergraduate Texts in Mathematics. Springer, Berlin, 2007.
- [8] F. R. Bernhart. Catalan, Motzkin, and Riordan numbers, *Discrete Math.* **204** (1999), 73–112.
- [9] A. Blecher, C. Brennan, and A. Knopfmacher. Combinatorial parameters in bargraphs, *Quaest. Math.* **39** (2016), 619–635.
- [10] A. Blecher, C. Brennan, and A. Knopfmacher. The site-perimeter of compositions, *Discrete Math. Appl.* **32** (2) (2022), 75–89.
- [11] A. Blecher, C. Brennan, A. Knopfmacher, and T. Mansour. The perimeter of words, *Discrete Math.* **340** (10) (2017), 2456–2465.
- [12] M. Bousquet-Mélou and A. Rechnitzer. The site-perimeter of bargraphs, *Adv. in Appl. Math.* **31**(2003), 86–112.
- [13] D. Callan, T. Mansour, and J. L. Ramírez. Statistics on bargraphs of Catalan words, *J. Autom. Lang. Comb.* **26** (2021), 177–196.
- [14] R. Donaghey and L. W. Shapiro. Motzkin numbers, *J. Combin. Theory Ser. A* **23**(3) (1977), 291–301.
- [15] P. Flajolet and R. Sedgewick. *Analytic Combinatorics*, Cambridge University Press, 2009.
- [16] I. M. Gessel and S. Ree. Lattice paths and Faber polynomials, *Advances in Combinatorial Methods and Applications to Probability and Statistics*, Birkhauser Verlag, Boston, 1997.
- [17] T. Goy and M. Shattuck. Determinant identities for the Catalan, Motzkin and Schröder numbers, *Art Disc. Appl. Math.* **7** (2024), #P1.09.
- [18] A. J. Guttmann (Ed.) *Polygons, Polyominoes and Polycubes*, Lecture Notes in Physics 775. Springer, Heidelberg, Germany, 2009.
- [19] C. Krattenthaler. Lattice path enumeration, In: Bóna, M. (ed.) *Handbook of Enumerative Combinatorics*, CRC Press, (2015).
- [20] T. Mansour. Semi-perimeter and inner site-perimeter of  $k$ -ary words and bargraphs, *Art Disc. Appl. Math.* **4** (2021), Article P1.06.
- [21] T. Mansour. The perimeter and the site-perimeter of set partitions, *Electron. J. Comb.* **26** (2) (2019), Article #P2.30.
- [22] T. Mansour. Interior vertices in set partitions, *Adv. in Applied Math.* **101** (2018), 60–69.
- [23] T. Mansour and A. Sh. Shabani. Enumerations on bargraphs, *Discrete Math. Lett.* **2** (2019), 65–94.
- [24] T. Mansour and J. L. Ramírez. Enumerations on polyominoes determined by Fuss-Catalan words, *Australas. J. Comb.* **81** (2021), 447–457.
- [25] T. Mansour and J. L. Ramírez. Exterior corners on bargraphs of Motzkin words, In: F. Hoffman, S. Holliday, Z. Rosen, F. Shahrokhi, and J. Wierman (eds). *Combinatorics, Graph Theory and Computing 2021*. Springer Proceedings in Mathematics & Statistics, vol 448. Springer, (2024).
- [26] T. Mansour, J. L. Ramírez, and D. A. Toquica. Counting lattice points on bargraphs of Catalan words, *Math. Comput. Sci.* **15** (2021), 701–713.
- [27] A. G. Orlov. On asymptotic behavior of the Taylor coefficients of algebraic functions, *Sib. Math. J.* **25**(5) (1994), 1002–1013.
- [28] H. Prodinger. The kernel method: a collection of examples, *Sém. Lothar. Combin.* **50** (2004), Paper B50f.
- [29] N. J. A. Sloane. The On-Line Encyclopedia of Integer Sequences, <http://oeis.org/>.
- [30] R. P. Stanley. *Catalan Numbers*, Cambridge University Press, 2015.

LIB, UNIVERSITÉ DE BOURGOGNE FRANCHE-COMTÉ, B.P. 47 870, 21078, DIJON CEDEX, FRANCE  
*Email address:* `barjl@u-bourgogne.fr`

LIB, UNIVERSITÉ DE BOURGOGNE FRANCHE-COMTÉ, B.P. 47 870, 21078, DIJON CEDEX, FRANCE  
*Email address:* `sergey.Kirgizov@u-bourgogne.fr`

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD NACIONAL DE COLOMBIA, BOGOTÁ, COLOMBIA  
*Email address:* `jlr Ramirez@unal.edu.co`

ESCUELA DE CIENCIAS EXACTAS E INGENIERÍA, UNIVERSIDAD SERGIO ARBOLEDA, BOGOTÁ, COLOMBIA  
*Email address:* `diego.villamizarr@usa.edu.co`